

**ADVANCED GCE
MATHEMATICS (MEI)**

Mechanics 3

4763

Candidates answer on the answer booklet.

OCR supplied materials:

- 8 page answer booklet (sent with general stationery)
- MEI Examination Formulae and Tables (MF2)

Other materials required:

- Scientific or graphical calculator

**Wednesday 22 June 2011
Morning**

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet. Please write clearly and in capital letters.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is being used.
- The total number of marks for this paper is **72**.
- This document consists of **8** pages. Any blank pages are indicated.

- 1 A particle is moving in a straight line. At time t its displacement x from a fixed point O on the line is given by

$$x = A \sin \omega t$$

where A and ω are constants.

(i) Show that $\frac{d^2x}{dt^2} = -\omega^2x$ and $\left(\frac{dx}{dt}\right)^2 = \omega^2(A^2 - x^2)$. [5]

A ball floats on the surface of the sea. Waves cause the ball to rise and fall in a vertical line, and the ball is executing simple harmonic motion. The centre of the oscillations is 8 m above the sea-bed. The ball has speed 1.2 m s^{-1} when it is 7.3 m above the sea-bed, and it has speed 0.75 m s^{-1} when it is 10 m above the sea-bed.

(ii) Show that the amplitude of the oscillations is 2.5 m, and find the period. [6]

(iii) Find the maximum speed of the ball. [1]

(iv) Find the magnitude and direction of the acceleration of the ball when it is 6.4 m above the sea-bed. [3]

(v) Find the time taken for the ball to move upwards from 6 m above the sea-bed to 9 m above the sea-bed. [3]

- 2 (a) A particle P of mass 0.2 kg is connected to a fixed point O by a light inextensible string of length 3.2 m, and is moving in a vertical circle with centre O and radius 3.2 m. Air resistance may be neglected. When P is at the highest point of the circle, the tension in the string is 0.6 N.
- (i) Find the speed of P when it is at the highest point. [3]
- (ii) For an instant when OP makes an angle of 60° with the downward vertical, find
- (A) the radial and tangential components of the acceleration of P, [5]
- (B) the tension in the string. [3]
- (b) A solid cone is fixed with its axis of symmetry vertical and its vertex V uppermost. The semi-vertical angle of the cone is 36° , and its surface is smooth. A particle Q of mass 0.2 kg is connected to V by a light inextensible string, and Q moves in a horizontal circle at constant speed, in contact with the surface of the cone, as shown in Fig. 2.




Fig. 2

The particle Q makes one complete revolution in 1.8 s, and the normal reaction of the cone on Q has magnitude 0.75 N.

- (i) Find the tension in the string. [2]
- (ii) Find the length of the string. [5]

- 3 Fixed points A and B are 4.8 m apart on the same horizontal level. The midpoint of AB is M. A light elastic string, with natural length 3.9 m and modulus of elasticity 573.3 N, has one end attached to A and the other end attached to B.

(i) Find the elastic energy stored in the string. [2]

A particle P is attached to the midpoint of the string, and is released from rest at M. It comes instantaneously to rest when P is 1.8 m vertically below M.

(ii) Show that the mass of P is 15 kg. [5]

(iii) Verify that P can rest in equilibrium when it is 1.0 m vertically below M. [4]

In general, a light elastic string, with natural length a and modulus of elasticity λ , has its ends attached to fixed points which are a distance d apart on the same horizontal level. A particle of mass m is attached to the midpoint of the string, and in the equilibrium position each half of the string has length h , as shown in Fig. 3.




Fig. 3

When the particle makes small oscillations in a vertical line, the period of oscillation is given by the formula

$$\sqrt{\frac{8\pi^2 h^3}{8h^3 - ad^2}} m^\alpha a^\beta \lambda^\gamma.$$

(iv) Show that $\frac{8\pi^2 h^3}{8h^3 - ad^2}$ is dimensionless. [1]

(v) Use dimensional analysis to find α , β and γ . [4]

(vi) Hence find the period when the particle P makes small oscillations in a vertical line centred on the position of equilibrium given in part (iii). [2]

- 4 The region A is bounded by the curve $y = x^2 + 5$ for $0 \leq x \leq 3$, the x -axis, the y -axis and the line $x = 3$. The region B is bounded by the curve $y = x^2 + 5$ for $0 \leq x \leq 3$, the y -axis and the line $y = 14$. These regions are shown in Fig. 4.




Fig. 4

- (i) Find the coordinates of the centre of mass of a uniform lamina occupying the region A . [9]
- (ii) The region B is rotated through 2π radians about the y -axis to form a uniform solid of revolution R . Find the y -coordinate of the centre of mass of the solid R . [6]
- (iii) The region A is rotated through 2π radians about the y -axis to form a uniform solid of revolution S . Using your answer to part (ii), or otherwise, find the y -coordinate of the centre of mass of the solid S . [3]

BLANK PAGE

BLANK PAGE

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.